

C R E D I T • V A L L E Y
THE CREDIT VALLEY HOSPITAL
PHYSICIAN'S ORDERS
PANTOPRAZOLE IV TO PO CONVERSION (Adult)

Weight _____(kg)
 Allergies _____

**To complete the order form, fill in the required blanks and/or check the appropriate boxes.
 To delete orders, draw one line through the item and initial.**

- | | | |
|---|-------|----------------------------------|
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV daily | START | Pantoprazole 40 mg PO daily* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV bid/Q12H | START | Pantoprazole 40 mg PO bid/Q12H* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV daily | START | Lansoprazole 30 mg NG daily* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV bid/Q12H | START | Lansoprazole 30 mg NG bid/Q12H** |

* **OR** automatic substitution to current Formulary Equivalent

As per IV-PO Conversion Policy approved at P&T and MAC

Inclusion Criteria

- Patient is tolerating other oral/NG medications x 24 hours

OR

- Patient is tolerating at least clear fluid diet x 24 hours

OR

- Patient is tolerating enteral feeds at goal rate x 24 hours

Date: _____ Time: _____

Pharmacist Signature



PHYSICIAN'S ORDERS

PANTOPRAZOLE IV TO PO CONVERSION (Adult)

Proton Pump Inhibitors: Pantoprazole IV to Pantoprazole PO/Lansoprazole NG Conversion

Indication for Use:

IV-PO Conversion:	Monitoring Only: Requires Physician Order for Conversion
Patients receiving IV Pantoprazole daily/BID because they are strictly NPO	Upper GI bleed prior to endoscopy in patients with: <ul style="list-style-type: none"> - Signs/symptoms of hemodynamic instability - Hypotension, postural hypotension or tachycardia - Active bleeding (hematemesis, hematochezia) - Upper GI bleed after endoscopic treatment in patients at high risk for re-bleeding Total days IV Pantoprazole Infusion: _____
Dose: Pantoprazole 40 mg IV daily or BID	Dose: Pantoprazole 80 mg loading dose followed by 8mg/h infusion

Dose Conversion:

- Pantoprazole 40 mg IV daily -> Pantoprazole 40 mg PO daily or Lansoprazole 30 mg NG daily
 - Pantoprazole 40 mg IV BID -> Pantoprazole 40 mg PO BID or Lansoprazole 30 mg NG BID
- OR current formulary equivalent

Inclusion Criteria:

- tolerating other oral/NG meds x 24 hours
- OR
- tolerating at least clear fluid diet x 24 hours
- OR
- tolerating enteral feeds at goal rate x 24 hours

Exclusion Criteria:

- NPO order on chart
- AND
- all medications administered by non-oral routes
- AND
- no oral/enteral feed intake

Reasons for no oral/enteral medication or feed intake:

- risk of aspiration (decreased LOC, seizures)
- GI obstruction
- complete bowel rest (inflammatory bowel disease, acute pancreatitis, fistula)
- pre/post-op fast
- nausea/vomiting/diarrhea
- continuous NG suction
- motility disorders of stomach/esophagus
- hemoglobin unstable
- high output ileostomy

Monitor for 48 hours to ensure that oral Pantoprazole is tolerated.



C R E D I T • V A L L E Y
THE CREDIT VALLEY HOSPITAL
PHYSICIAN'S ORDERS
PANTOPRAZOLE IV TO PO CONVERSION (Adult)

Weight _____(kg)
 Allergies _____

**To complete the order form, fill in the required blanks and/or check the appropriate boxes.
 To delete orders, draw one line through the item and initial.**

- | | | |
|---|-------|----------------------------------|
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV daily | START | Pantoprazole 40 mg PO daily* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV bid/Q12H | START | Pantoprazole 40 mg PO bid/Q12H* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV daily | START | Lansoprazole 30 mg NG daily* |
| <input type="checkbox"/> Discontinue Pantoprazole 40 mg IV bid/Q12H | START | Lansoprazole 30 mg NG bid/Q12H** |

* **OR** automatic substitution to current Formulary Equivalent

As per IV-PO Conversion Policy approved at P&T and MAC

Inclusion Criteria

- Patient is tolerating other oral/NG medications x 24 hours

OR

- Patient is tolerating at least clear fluid diet x 24 hours

OR

- Patient is tolerating enteral feeds at goal rate x 24 hours

Date: _____ **Time:** _____

Pharmacist Signature



PHYSICIAN'S ORDERS

PANTOPRAZOLE IV TO PO CONVERSION (Adult)

Proton Pump Inhibitors: Pantoprazole IV to Pantoprazole PO/Lansoprazole NG Conversion

Indication for Use:

IV-PO Conversion:	Monitoring Only: Requires Physician Order for Conversion
Patients receiving IV Pantoprazole daily/BID because they are strictly NPO	Upper GI bleed prior to endoscopy in patients with: <ul style="list-style-type: none"> - Signs/symptoms of hemodynamic instability - Hypotension, postural hypotension or tachycardia - Active bleeding (hematemesis, hematochezia) - Upper GI bleed after endoscopic treatment in patients at high risk for re-bleeding Total days IV Pantoprazole Infusion: _____
Dose: Pantoprazole 40 mg IV daily or BID	Dose: Pantoprazole 80 mg loading dose followed by 8mg/h infusion

Dose Conversion:

- Pantoprazole 40 mg IV daily -> Pantoprazole 40 mg PO daily or Lansoprazole 30 mg NG daily
 - Pantoprazole 40 mg IV BID -> Pantoprazole 40 mg PO BID or Lansoprazole 30 mg NG BID
- OR current formulary equivalent

Inclusion Criteria:

- tolerating other oral/NG meds x 24 hours
- OR
- tolerating at least clear fluid diet x 24 hours
- OR
- tolerating enteral feeds at goal rate x 24 hours

Exclusion Criteria:

- NPO order on chart
- AND
- all medications administered by non-oral routes
- AND
- no oral/enteral feed intake

Reasons for no oral/enteral medication or feed intake:

- risk of aspiration (decreased LOC, seizures)
- GI obstruction
- complete bowel rest (inflammatory bowel disease, acute pancreatitis, fistula)
- pre/post-op fast
- nausea/vomiting/diarrhea
- continuous NG suction
- motility disorders of stomach/esophagus
- hemoglobin unstable
- high output ileostomy

Monitor for 48 hours to ensure that oral Pantoprazole is tolerated.

